Re-evaluating Crop Nutrient Management In Light Of Spatial Variability: Orchard Crops

Patrick Brown University of California, Davis

What do we know and how do we manage? Leaf Sampling and Critical Value Analysis in Orchard crops

(based on Ulrich and Hills @ UC Davis in 1950-70's)

 Table 26.2
 Critical nutrient levels (dry-weight basis) in almond leaves sampled in July.



*Critical values for boron deficiency and toxicity are currently being revised. Hull boron >300 ppm is excessive. Leaf sampling is not effective to determine excess boron.

Composite leaf samples from well defined locations in tree

- SW corner, non-fruiting, fully exposed leaves
- July/August values only
- Typically 1 composite sample per management unit

Existing Standards were based on

Limited field/yield trials (N, K, B) or,

Appearance of symptoms (P, S, Mg, Ca, Mn, Zn, Fe, Cu) or.

- ➤Unknown (Ni, Cl, Mo)
- ≻Experience

Almond Grower Survey 2008

Patrick Brown, Cary Trexler, Sara Lopus, Maria Santibanez

How well are current nutrient management guidelines understood, utilized and implemented?

- Focus Group Activity
 - 45 leading Growers, Consultants and University Representatives (FA, Extension, Govt.)
- Random, balanced selection of 1,650 Almond Growers
 - 558 responses (33% of industry)
 - adequate to ensure +/- 5% margin of error

Almond is a \$2 billion crop and California's largest export crop.



Do you think the University of California critical values are adequate to ensure maximal productivity in almonds?



Apparently we are not as effective as we should be - Why?

Is the use of Plant Samples and the Critical Value or Critical Range appropriate for Trees/Vines?

- Early German agronomic work (1890's), US development and adaptation (Ulrich & Hills - UCD) applied to trees at UC Davis and elsewhere (1950's - present) and used worldwide with only modest new investigation.
- In Agronomic crops, detecting and correcting deficiencies has historically been the central principle
 - Soil is depleted until a response is seen.
 - In trees, prevention, not correction is the goal of a good manager.
- Establishment of CV's and monitoring nutrient status in trees is complex.
 - Multi-year relationship between nutrition and yield spans many years
 - Distribution of nutrients in the perennial crops and fields is extremely variable.
 - Adequate sampling is difficult.
- Problems with implementation and interpretation (known, unknown and forgotten)
 - Incomplete and inadequate information.
 - Lack of ability to translate results to actions.
 - Realities of orchard design and management practice.

Effect of K on Yield in Almond

Nutrient Interactions





Standard Sample: Fully Exposed non-fruiting leaves in late summer

Strong Yield Interactions High Nutrition is essential for High Yield High Yield however depresses Leaf Nutrients Multi-year Nutrient x Yield Correlations are Strikingly Difficult to Interpret.

Leaf Analysis: Problems and Challenges:

•Recommended protocols and CV's are 'soft'.

•Tissue sampling and yield determination errors compromise

utility of tissue sampling.

Confounding Factors (multi-element deficiencies, drought, disease)

• Clear interdependency between yield, nutrient demand and nutrient status; yield/nutrition and potential yield.

• Non linear, variable, multi-year relationship between yield and nutrient response.



A.K. Alva et al. / Scientia Horticulturae 107 (2006) 233-244

The Most Immediate Problem With Our Use of Critical Values





Pistachio Yield



K Variability and Optimization in Almond

Leaf samples collected from 50 tree rows.

Grower target CV = 2.0% K (95% of trees are above 1.4%K)

University of California recommended CV = 1.4% K) On a purely economic basis growers are clearly making the right choice though they are doing so with:-

No knowledge of the rationale behind this practice
No recognition of the constraints of tissue sampling and that field mean leaf nutrient concentrations cannot be interpreted without knowledge of the variability in the field.

No knowledge of nutrient response dynamics or returns

•No consideration of the environmental consequence of their practices.

Alternatives?



Alternate Approaches to Nutrient Management in Trees

Nutrient Budgets (EU Model) I=D x E

Replacing nutrients removed from the orchard or vineyard, minimizing non-crop

export.

Essential Components and Challenges:

- Demand
 - Annual Demand and Variability
 - Seasonal patterns of demand and uptake
- Inputs and losses
 - Fertilizer
 - Irrigation
 - Soil Mineralization (timing and quantity, environment and management interactions)
 - Storage in perennial tissues
 - Leaching, Volatilization
 - Cover Crops/Manures, Atmospheric Deposition, Crop residues.
- Efficiencies and Interactions
 - Synchronization and synlocation
 - Source and method of applications
 - Other determinants

Nutrient Demand: Whole tree Harvesting: 5 mature trees x 5 times in a year

Whole Tree N Contents by Organ in Almond.



Almond Variation within orchard

Fresh weights





Yield is not uniform in any field. Yield of 10,040 trees Pistachio trees (40 ha)





Pistachio Yield



Managing for Annual Yield N.

Requires Yield Estimate (April) or Yield Model





Pistachio Yield



Influence of Precision Management of Fertilizer Losses – first steps.



■ 2002 ■ 2003 ■ 2004 ■ 2005 ■ 2006 **■** 2007

Pattern Recognition and Yield Estimation in Pistachio.



Whole field yield has been successfully modeled (+/- 30%) based on:

Historic yieldClimate

Chilling hours, heat units, weather anomalies.
Early season predictors
Sampling procedures
Remote sensing

Individual tree determination remains more challenging: •Sub populations of trees clearly exist •Biological basis for yield fluctuation is not well understood.

Chaos Dynamics, non linear modeling and the Prediction of Yield in Satsuma Mandarin

Kenshi Sakai, Tokyo University of Agriculture and Technology

Off year tree

On year tree





Estimation of Jaccobian dynamics from ensemble data set of 96 individuals over 5 years.



One Year Forward Prediction



Model does not utilize any biological principles or environmental variables.

Pistachio data set is 10,000 individuals for 6 years with information on plant biology and environment.

30

predicted

Almond trial initiated 2008. 7,000 individuals at 4 sites with extensive measurement of biological and environmental variables.

Yield prediction is possible.

000000000



Orchard yield variability compromises research and inference

Statistical power and model assumptions



•In off-years yield is not normally distributed.

•There is significant auto-correlation within and between years and the degree of auto-correlation varies with yield.

•These two observations essentially invalidate traditional statistical procedures.

Testing Experimental Designs for Orchard Research

Design id.	Experimental design	Blocks (#)	Block size	Reps. per block	Reps. per treat.	Total N	Description
1	Completely random	-	-	-	27	81	No predetermined spatial configuratioin
2	Large blocks random	3	6 x 6	9	27	81	27 randomly selected out of 36 in each block
3	Small rectangle not replicated	27	3 x 1	1	27	81	0 reps of 3 randomly assigned treatments over 27 blocks
4	Randomized within row	9	9 x 1	3	27	81	3 reps of 3 treatments randomly assigned in each of 9 rows
5	Random row	9	-	9	27	81	9 reps of 1 treatment in each of 9 rows
6	Medium square	3	5 x 5	3	27	81	same as design #7 except an unused unit between every treatment
7	Small square	3	3 x 3	3	27	81	3 reps of 3 treatments randomly assigned within a square
8	Grid- based entire	_	-	_	27	81	81 randomly assigned treatment equally spaced over entire orchard
9	Grid-based half	-	-	_	27	81	81 randomly assigned treatments equally spaced over east or west half
10	Grid based quadrants	-	-	-	27	81	81 randomly assigned treatments equally spaced over one quadrants

Testing Experimental Designs for Orchard Research



Experimental design id.

Nutrient Use Efficiency of 4820 Individual Tree over 6 Years.

(Fertilizer application per tree / Nutrient Export per tree (nuts)



Re-evaluating Crop Nutrient Management In Light Of Spatial Variability in Orchard Crops

The high value and long life of perennial systems, the inadequacy of current practices, the willingness of industry to adopt technology and above all, the environmental and market demands for better management practices, represents an ideal opportunity to re-examine and re-invent our approach to nutrient management in high value crops.

This Requires:

•Yield Measurement and Prediction – Integrated mathematical, biological, engineering and ecological approaches.

•Determination of Spatial Variability - Statistical and geo-statistical tools, sampling and sensing technologies, improved experimental designs.

•Improved Fertilizer Efficiency - Agronomic and physiological experimentation to optimize rates, timing, formulation.

•New Management Tools – Rapid yield and nutrient measurement techniques. New approaches to precision application -sub sector fertigation to single tree fertigation; VR devices and materials (surface/liquid)

•Design and extension of flexible and easy to use decision support systems.

XVI International Plant Nutrition Colloquium



"Plant Nutrition for Sustainable Development and Global Health"

Sacramento, California, USA

Aug 26-30, 2009

Registration, call for papers and program available at

<u>ipnc.ucdavis.edu</u>

Hosted by the **Department of Plant Sciences** and the University of California, Davis

